

## Weight-Loss Surgery HRA v3

### Client Review Document

#### Overview

The Weight-Loss Surgery HRA uses body mass index (BMI), quality of life (QoL), and health history to assess possible eligibility for weight-loss surgery.<sup>1,2</sup> It also calculates the number of pounds away from the healthy weight range (BMI = 24.9) for overweight users and gives a weight-loss recommendation.<sup>3,4</sup>

#### Main Scientific Basis

This HRA was designed using the 2013 clinical practice guidelines cosponsored by the American Association of Clinical Endocrinologists (AACE), The Obesity Society (TOS), and the American Society for Metabolic & Bariatric Surgery (ASMBS).<sup>1</sup>

#### Product Description

The Weight-Loss Surgery HRA asks a series of health history questions and uses clinical practice guidelines to identify possible bariatric surgery candidates.<sup>1,2</sup> The HRA also makes a weight loss recommendation irrespective of surgery eligibility.<sup>3,4</sup> Several weight-related health conditions are identified, including: hypertension, abnormal cholesterol, prediabetes or type 2 diabetes, arthritis, asthma, obstructive sleep apnea, gastroesophageal reflux disease, urinary incontinence, obesity-hypoventilation syndrome, pseudotumor cerebri, nonalcoholic liver disease, and venous stasis disease.

#### Key Results Provided

The primary result from the Weight-Loss Surgery HRA is eligibility for bariatric surgery. Results are categorized as *possible surgery candidate*, *not a surgery candidate*, and *screening needed to determine eligibility*. Follow-up messaging, emails, and programs can be developed to align with these categories.

#### About Bariatric Surgery Eligibility

In the following 3 cases, bariatric surgery may be recommended: class 3 obesity (BMI = 40 or higher); class 2 obesity (BMI = 35.0-39.9) with a weight-related health condition or greatly impaired QoL;<sup>1</sup> class 1 obesity (BMI = 30.0-34.9) with uncontrolled type 2 diabetes.<sup>2</sup> People with class 2 obesity who don't answer enough questions to determine surgery eligibility are referred for follow-up.

## About Weight Loss Recommendation

Weight loss is recommended for people with obesity (BMI = 30 or higher) and for people with overweight (BMI = 25.0-29.9) who have a weight-related health condition or increased waist circumference.<sup>3-4</sup> Overweight people with no weight-related health conditions are told to avoid weight gain. Those who don't answer enough questions to determine a weight-loss recommendation are referred for follow-up.

## The Role of QoL

The impact of weight on QoL is used in the bariatric surgery algorithm but not in the weight loss recommendation. Regardless of surgery eligibility or weight-loss recommendation, people who report that their weight greatly impacts their QoL are urged to talk to a health care professional about their weight.

## References

1. Mechanick JI, Youdim A, Jones DB, et al. Clinical Practice Guidelines for the Perioperative Nutritional, Metabolic, and Nonsurgical Support of the Bariatric Surgery Patient—2013 Update: Cosponsored by American Association of Clinical Endocrinologists, The Obesity Society, and American Society for Metabolic & Bariatric Surgery. *Surg Obes Relat Dis*. 2013;9:159–191. DOI: <https://doi.org/10.1016/j.soard.2012.12.010>
2. Rubino F, Nathan DM, Eckel RH, et al. Metabolic surgery in the treatment algorithm for type 2 diabetes: a joint statement by international diabetes organizations. *Diabetes Care*. 2016;39:861-877. DOI: <https://doi.org/10.2337/dc16-0236>
3. Jensen MD, Ryan DH, Apovian CM, Ard JD, Comuzzie AG, Donato KA, et. al. 2013 AHA/ACC/TOS Guideline for the Management of Overweight and Obesity in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Circulation*. 2013;00.000-000. DOI: <http://dx.doi.org/10.1161/01.cir.0000437739.71477.ee>
4. Grundy SM, Cleeman JI, Daniels SR, et al. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation*. 2005;112:2735-2752. DOI: <https://doi.org/10.1161/CIRCULATIONAHA.105.169404>